

sequences provided herein are based on known sequencing techniques.

Analysis of the amino acid sequence of the full-length PRO1008 polypeptide suggests that portions of it possess significant sequence identity and/or similarity with mdkk-1, thereby indicating that PRO1008 may be a novel member of this family and have head inducing activity.

Still analyzing the amino acid sequence of SEQ ID NO:456, the putative signal peptide is at about amino acids 1-23 of SEQ ID NO:456. The N-glycosylation site is at about amino acids 256-259 of SEQ ID NO:456, and the fungal zn-(2)-cys(6) binuclear cluster domain is at about amino acids 110-126 of SEQ ID NO:456. The corresponding nucleotides can of all the amino acids can be routinely determined given the sequences provided herein.

EXAMPLE 76: Isolation of cDNA Clones Encoding Human PRO1012

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above, wherein the consensus sequence is herein designated DNA49313. Based on the DNA49313 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1012.

A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-ACTCCCCAGGCTGTTCACACTGCC-3' (SEQ ID NO:460);

reverse PCR primer 5'-GATCAGCCAGCCAATACCAGCAGC-3' (SEQ ID NO:461).

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the DNA49313 consensus sequence which had the following nucleotide sequence:

hybridization probe

5'-GTGGTGATGATAGAATGCTTTGCCGAATGAAAGGAGTCAACAGCTATCCC-3' (SEQ ID NO:462).

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1012 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue (LIB227).

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1012 [herein designated as UNQ495 (DNA56439-1376)] (SEQ ID NO:458) and the derived protein sequence for PRO1012.

The entire nucleotide sequence of UNQ495 (DNA56439-1376) is shown in Figure 189 (SEQ ID NO:458). Clone UNQ495 (DNA56439-1376) contains a single open reading frame with an apparent translational initiation site at nucleotide positions 404-406 and ending at the stop codon at nucleotide positions 2645-2647 (Figure 189). The predicted polypeptide precursor is 747 amino acids long (Figure 190). The full-length PRO1012 protein shown in Figure 190 has an estimated molecular weight of about 86,127 daltons and a pI of about 7.46. Clone UNQ495 (DNA56439-1376) has been deposited with ATCC on May 14, 1998. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

Analysis of the amino acid sequence of the full-length PRO1012 polypeptide suggests that portions of it possess sequence identity with disulfide isomerase thereby indicating that PRO1012 may be a novel disulfide isomerase related protein.

Still analyzing the amino acid sequence of SEQ ID NO:459, the cytochrome C family heme-binding site signature is at about amino acids 158-163 of SEQ ID NO:459. The Nt-DNAJ domain signature is at about amino acids 77-96 of SEQ ID NO:459. An N-glycosylation site is at about amino acids 484-487 of SEQ ID NO:459.

The ER targeting sequence is at about amino acids 744-747 of SEQ ID NO:459. It is understood that the polypeptide and nucleic acids disclosed can be routinely formed with or without, these portions as desired, in alternative embodiments. For example, it may be desirable to produce PRO1012 without the ER targeting sequence. The corresponding nucleotides can be routinely determined given the sequences provided herein.

EXAMPLE 77: Isolation of cDNA Clones Encoding Human PRO1014

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above, wherein the consensus sequence obtained is herein designated DNA49811. Based upon an observed homology between the DNA49811 sequence and Incyte EST clone no. 2612207, Incyte EST clone no. 2612207 was purchased and its insert was obtained and sequenced, wherein the sequence obtained is shown in Figure 191 (SEQ ID NO:463).

DNA sequencing gave the full-length DNA sequence for PRO1014 [herein designated as UNQ497 (DNA56409-1377)] (SEQ ID NO:463) and the derived protein sequence for PRO1014.

The entire nucleotide sequence of UNQ497 (DNA56409-1377) is shown in Figure 191 (SEQ ID NO:463). Clone UNQ497 (DNA56409-1377) contains a single open reading frame with an apparent translational initiation site at nucleotide positions 66-68 and ending at the stop codon at nucleotide positions 966-968 (Figure 191). The predicted polypeptide precursor is 300 amino acids long (Figure 192). The full-length PRO1014 protein shown in Figure 192 has an estimated molecular weight of about 33,655 daltons and a pI of about 9.31. Clone UNQ497 (DNA56409-1377) has been deposited with the ATCC on May 20, 1998. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

Analysis of the amino acid sequence of the full-length PRO1014 polypeptide suggests that portions of it possess sequence identity with reductase, thereby indicating that PRO1014 may be a novel member of the reductase family.

Still analyzing the amino acid sequence of SEQ ID NO:464, the putative signal peptide is at about amino acids 1-19 of SEQ ID NO:464. The cAMP and cGMP dependent protein kinase phosphorylation sites are at about amino acids 30-33 and 58-61 of SEQ ID NO:464. Short chain alcohol dehydrogenase family proteins are at about amino acids 165-202, 37-49, 112-122 and 210-219 of SEQ ID NO:464. The corresponding nucleotides of these domains and any other amino acids provided herein can be routinely determined given the sequences provided herein.

EXAMPLE 78: Isolation of cDNA Clones Encoding Human PRO1017

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described

in Example 1 above, wherein that consensus DNA sequence is herein designated DNA53235. Based upon an observed homology between the DNA53235 consensus sequence and the Merck EST clone no. AA243086, the Merck EST clone no. AA243086 was purchased and its insert obtained and sequenced, wherein the sequence obtained is shown in Figure 193 (SEQ ID NO:465). DNA sequencing gave the full-length DNA sequence for PRO1017 [herein designated as UNQ500 (DNA56112-1379)] (SEQ ID NO:465) and the derived protein sequence for PRO1017.

The entire nucleotide sequence of UNQ500 (DNA56112-1379) is shown in Figure 193 (SEQ ID NO:465). Clone UNQ500 (DNA56112-1379) contains a single open reading frame with an apparent translational initiation site at nucleotide positions 128-130 and ending at the stop codon at nucleotide positions 1370-1372 (Figure 193). The predicted polypeptide precursor is 414 amino acids long (Figure 194). The full-length PRO1017 protein shown in Figure 194 has an estimated molecular weight of about 48,414 daltons and a pI of about 9.54. Clone UNQ500 (DNA56112-1379) has been deposited with the ATCC. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

Analysis of the amino acid sequence of the full-length PRO1017 polypeptide suggests that portions of it possess sequence identity with HNK-1 sulfotransferase, thereby indicating that PRO1017 may be a novel sulfotransferase.

Still analyzing the amino acid sequence of SEQ ID NO:466, the putative signal peptide is at about amino acids 1-31 of SEQ ID NO:466. N-glycosylation sites are at about amino acids 134-137, 209-212, 280-283 and 370-273 of SEQ ID NO:466. The TNFR/NGFR family cystein-rich region protein is at about amino acids 329-332 of SEQ ID NO:466. The corresponding nucleotides can be routinely determined given the sequences provided herein. The protein can be secreted.

EXAMPLE 79: Isolation of cDNA Clones Encoding Human PRO474

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above, wherein the consensus sequence obtained is herein designated DNA49818. Based upon an observed homology between the DNA49818 consensus sequence and the Merck EST clone no. H77889, the Merck EST clone no. H77889 was purchased and its insert obtained and sequenced, wherein the sequence obtained is herein shown in Figure 195 (SEQ ID NO:467). DNA sequencing gave the full-length DNA sequence for PRO474 [herein designated as UNQ502 (DNA56045-1380)] (SEQ ID NO:467) and the derived protein sequence for PRO474.

The entire nucleotide sequence of UNQ502 (DNA56045-1380) is shown in Figure 195 (SEQ ID NO:467). Clone UNQ502 (DNA56045-1380) contains a single open reading frame with an apparent translational initiation site at nucleotide positions 106-108 and ending at the stop codon at nucleotide positions 916-918 (Figure 195). The predicted polypeptide precursor is 270 amino acids long (Figure 196). The full-length PRO474 protein shown in Figure 196 has an estimated molecular weight of about 28,317 daltons and a pI of about 6.0. Clone UNQ502 (DNA56045-1380) has been deposited with the ATCC. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.